

5
sup
31

~~tion i
3, fil
ted in~~

- 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100 101 102 103 104 105 106 107 108 109 110 111 112 113 114 115 116 117 118 119 120 121 122 123 124 125 126 127 128 129 130 131 132 133 134 135 136 137 138 139 140 141 142 143 144 145 146 147 148 149 150 151 152 153 154 155 156 157 158 159 160 161 162 163 164 165 166 167 168 169 170 171 172 173 174 175 176 177 178 179 180 181 182 183 184 185 186 187 188 189 190 191 192 193 194 195 196 197 198 199 200 201 202 203 204 205 206 207 208 209 210 211 212 213 214 215 216 217 218 219 220 221 222 223 224 225 226 227 228 229 230 231 232 233 234 235 236 237 238 239 240 241 242 243 244 245 246 247 248 249 250 251 252 253 254 255 256 257 258 259 260 261 262 263 264 265 266 267 268 269 270 271 272 273 274 275 276 277 278 279 280 281 282 283 284 285 286 287 288 289 290 291 292 293 294 295 296 297 298 299 300 301 302 303 304 305 306 307 308 309 310 311 312 313 314 315 316 317 318 319 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 409 410 411 412 413 414 415 416 417 418 419 420 421 422 423 424 425 426 427 428 429 430 431 432 433 434 435 436 437 438 439 440 441 442 443 444 445 446 447 448 449 450 451 452 453 454 455 456 457 458 459 460 461 462 463 464 465 466 467 468 469 470 471 472 473 474 475 476 477 478 479 480 481 482 483 484 485 486 487 488 489 490 491 492 493 494 495 496 497 498 499 500 501 502 503 504 505 506 507 508 509 510 511 512 513 514 515 516 517 518 519 520 521 522 523 524 525 526 527 528 529 530 531 532 533 534 535 536 537 538 539 540 541 542 543 544 545 546 547 548 549 550 551 552 553 554 555 556 557 558 559 560 561 562 563 564 565 566 567 568 569 570 571 572 573 574 575 576 577 578 579 580 581 582 583 584 585 586 587 588 589 590 591 592 593 594 595 596 597 598 599 600 601 602 603 604 605 606 607 608 609 610 611 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 648 649 650 651 652 653 654 655 656 657 658 659 660 661 662 663 664 665 666 667 668 669 670 671 672 673 674 675 676 677 678 679 680 681 682 683 684 685 686 687 688 689 690 691 692 693 694 695 696 697 698 699 700 701 702 703 704 705 706 707 708 709 710 711 712 713 714 715 716 717 718 719 720 721 722 723 724 725 726 727 728 729 730 731 732 733 734 735 736 737 738 739 740 741 742 743 744 745 746 747 748 749 750 751 752 753 754 755 756 757 758 759 760 761 762 763 764 765 766 767 768 769 770 771 772 773 774 775 776 777 778 779 780 781 782 783 784 785 786 787 788 789 790 791 792 793 794 795 796 797 798 799 800 801 802 803 804 805 806 807 808 809 810 811 812 813 814 815 816 817 818 819 820 821 822 823 824 825 826 827 828 829 830 831 832 833 834 835 836 837 838 839 840 841 842 843 844 845 846 847 848 849 850 851 852 853 854 855 856 857 858 859 860 861 862 863 864 865 866 867 868 869 870 871 872 873 874 875 876 877 878 879 880 881 882 883 884 885 886 887 888 889 890 891 892 893 894 895 896 897 898 899 900 901 902 903 904 905 906 907 908 909 910 911 912 913 914 915 916 917 918 919 920 921 922 923 924 925 926 927 928 929 930 931 932 933 934 935 936 937 938 939 940 941 942 943 944 945 946 947 948 949 950 951 952 953 954 955 956 957 958 959 960 961 962 963 964 965 966 967 968 969 970 971 972 973 974 975 976 977 978 979 980 981 982 983 984 985 986 987 988 989 990 991 992 993 994 995 996 997 998 999 1000 1001 1002 1003 1004 1005 1006 1007 1008 1009 1010 1011 1012 1013 1014 1015 1016 1017 1018 1019 1020 1021 1022 1023 1024 1025 1026 1027 1028 1029 1030 1031 1032 1033 1034 1035 1036 1037 1038 1039 1040 1041 1042 1043 1044 1045 1046 1047 1048 1049 1050 1051 1052 1053

Background of the Invention

- 25 Cancer is a complex multifactor and multistep process
involving the coordinated expression and suppression of
genes functioning as positive and negative regulators of
oncogenesis (1-5). Direct cloning strategies, based on
transfer of a dominant transforming or tumorigenic
30 phenotype, have identified positive acting oncogenes
(6-9). In contrast, the detection and cloning of genes
that suppress the cancer phenotype have proven more
difficult and elusive (10-15). A direct approach for
isolating genes directly involved in regulating growth
35 and differentiation involves subtraction hybridization
between cDNA libraries constructed from actively growing
cancer cells and cDNA libraries from cancer cells induced
to lose proliferative capacity irreversibly and
terminally differentiate (13,14). This experimental
40 strategy has been applied to human melanoma cells,

- 2 -

induced to terminally differentiate by treatment with recombinant human interferon β (IFN- β) and mezerein (MEZ), resulting in the cloning of novel melanoma differentiation-associated (*mda*) genes not previously described in DNA data bases (13,14). A direct role for specific *mda* genes in mediating growth and cell cycle control is apparent by the identification and cloning of *mda-6* (13-16), which is identical to the ubiquitous inhibitor of cyclin-dependent kinases p21 (17). The importance of p21 in growth control is well documented and this gene has been independently isolated, as WAF-1, CIP-1, and SDI-1, by a number of laboratories using different approaches (18-20). These studies indicate that specific genes associated with proliferative control are induced and may contribute to the processes of growth arrest and terminal differentiation in human cancer cells.

The *mda-7* gene was cloned from a differentiation inducer (IFN- β plus MEZ)-treated human melanoma (H0-1) subtracted library (13,14). The full-length *mda-7* cDNA is 1718 nucleotides, and the major open reading frame encodes a novel protein of 206 aa with an M_r of 23.8 kDa (21). Previous studies indicate that *mda-7* is induced as a function of growth arrest and induction of terminal differentiation in human melanoma cells (14,21). *mda-7* expression also inversely correlates with melanoma progression - i.e., actively growing normal human melanocytes express more *mda-7* than metastatic human melanoma cells (21). Moreover, *mda-7* is growth inhibitory toward human melanoma cells in transient transfection assays and in stable transformed cells containing a dexamethasone (DEX)-inducible *mda-7* gene (21). These studies indicate that *mda-7* may contribute to the physiology of human melanocytes and melanomas, and this gene has growth suppressive properties when overexpressed in human melanoma cells.

- 3 -

The *mda-7* gene was also described in the International Patent Cooperation Treaty Application No. PCT/US94/12160, international filing date, October 24, 1994 with International Publication No. WO95/11986, the content of which is incorporated into this application by reference.

This invention reports that *mda-7* is a potent growth suppressing gene in cancer cells of diverse origin, including breast, central nervous system, cervix, colon, prostate and connective tissue. An inhibition in colony formation occurs in cancer cells containing defects in their p53 and/or retinoblastoma (RB) genes or lacking p53 and RB expression. In contrast, expression of *mda-7* in normal human mammary epithelial cells, human skin fibroblasts and rat embryo fibroblasts induces quantitatively less growth suppression than in cancer cells. When stably expressed in human cervical carcinoma (HeLa) and prostate carcinoma (DU-145) cells, *mda-7* has a negative effect on growth and transformation-related properties. The effects of *mda-7* on HeLa cells are reversible following abrogation of the MDA-7 protein by infection with a genetically modified Ad5 vector expressing an antisense *mda-7* gene. These observations indicate that *mda-7* is a novel growth suppressing gene with a wide range of inhibitory actions in human cancers manifesting different genetic defects.

09944345A

- 4 -

Summary of the Invention

5 This invention provides a method for reversing the cancerous phenotype of a cancer cell by introducing a nucleic acid including a melanoma differentiation associated gene (*mda-7*) into the cell under conditions permitting the expression of the gene so as to thereby reverse the cancerous phenotype of the cell. This invention also provides a method for reversing the
10 cancerous phenotype of cancer cell in a subject by introducing the above-described nucleic acid into the subject's cancerous cell.

15 This invention also provides a method for reversing the cancerous phenotype of a cancer cell by introducing the gene product of a melanoma differentiation associated gene (*mda-7*) into the cancer cell so as to thereby reverse the cancerous phenotype of the cell. This invention also provides a method for reversing the
20 cancerous phenotype of a cancer cell in a subject by introducing the above-described gene product into the subject's cancerous cell.

25 This invention also provides a pharmaceutical composition having an amount of a nucleic acid including a melanoma differentiation associated gene (*mda-7*) effective to reverse the cancerous phenotype of a cancer cell and a pharmaceutically acceptable carrier. This invention also provides a pharmaceutical composition having an amount of
30 the gene product of the above-described gene effective to reverse the cancerous phenotype of a cancer cell and a pharmaceutically acceptable carrier.

00004452-112104

- 5 -

Brief Description of the Figures

5 **Figure 1.** Effect of *mda-7* expression on hygromycin resistant colony formation in HeLa cells. HeLa cells were transfected with 10 µg of pREP4 vector (RSV-vector), *mda-7* cloned in an antisense orientation in the pREP4 vector (RSV-MDA-7-Antisense), or *mda-7* cloned in a sense orientation in the pREP4 vector (RSV-MDA-7-Sense) and selected in media
10 containing 100 µg of hygromycin.

15 **Figure 2.** Effect of antisense *mda-7* on monolayer growth of pREP4 vector HeLa cl 1 and *mda-7* (S) expressing HeLa cl 2 cells. HeLa cl 1 (pREP4 vector transformed HeLa clone) and HeLa cl 2 (*mda-7* expressing HeLa clone) cells were grown in the absence or following infection with 10 plaque forming units/cell with a recombinant type 5 adenovirus (Ad5) expressing antisense
20 *mda-7* [Ad.*mda-7* (AS)]. Results are the average cell number from triplicate samples that varied by ≤ 10%.

25 **Figure 3.** Effect of antisense *mda-7* on the high molecular weight-MDA-7 complexing (HMC) protein, the MDA-7 protein and the actin protein in HeLa, HeLa cl 1, and HeLa cl 2 cells. HeLa and HeLa cl 1 (pREP4 vector transformed HeLa clone) were uninfected (-) or infected (+) with 10 plaque forming units/cell of Ad.*mda-7* (AS) for 96 hr
30 labeled with [³⁵S] methionine, and the levels of the HMC, MDA-7 and actin proteins were determined by immunoprecipitation analysis. For HeLa cl 2 (*mda-7* expressing HeLa clone), the
35 effect of infection with 10 plaque forming units/ml of Ad.*mda-7* (AS) on protein levels was determined by immunoprecipitation analysis of

00991452.112104

- 6 -

[³⁵S] methionine labeled cell lysates after +24, +48, +72 and +96 hr. The effect of infection of HeLa cl 2 cells with the control mutant Ad5, H5dl 434, was determined by immunoprecipitation analysis of [³⁵S]methionine labeled cell lysates 96 hr after infection with 10 plaque forming units/cell.

Figures 4A & 4B Synthesis of *mda-7* RNA and protein in DU-145 clones containing a DEX-inducible *mda-7* gene.

Figure 4A. Cells were grown in the absence or presence of 10⁻⁶ M DEX for 96 hr, and total RNA was isolated, subjected to Northern blotting and probed with *mda-7*, a neomycin resistance (Neo^R) gene and GAPDH.

Figure 4B. Cells were grown in the absence or presence of 10⁻⁶ M DEX for 96 hr, cellular proteins were labeled with [³⁵S] methionine and immunoprecipitated with antibodies recognizing MDA-7 and actin proteins.

Figure 5 Inhibition of growth of established human cervical cancer (HeLa) xenografts in athymic nude mice.

Figure 6 Effect of Ad.*mda-7* S on HeLa tumor volume ratios. The result indicates that Ad.*mda-7* S can inhibit tumor progression in vivo in nude mice.

- 7 -

Detailed Description of the Invention

In order to facilitate an understanding of the Experimental Details section which follows, certain frequently occurring methods and/or terms are described in Sambrook, et al. (45).

This invention provides a method for reversing the cancerous phenotype of a cancer cell which comprises introducing a nucleic acid comprising a melanoma differentiation associated gene (*mda-7*) into the cell under conditions permitting the expression of the gene so as to thereby reverse the cancerous phenotype of the cell.

This invention also provides a method for reversing the cancerous phenotype of a cancer cell in a subject which comprises introducing a nucleic acid molecule comprising a melanoma differentiation associated gene (*mda-7*) into the subject's cancerous cell under conditions permitting expression of the gene in the subject's cells so as to thereby reverse the cancerous phenotype of the cell.

Methods to introduce a nucleic acid molecule into cells have been well known in the art. Naked nucleic acid molecule may be introduced into the cell by direct transformation. Alternatively, the nucleic acid molecule may be embedded in liposomes. Accordingly, this invention provides the above methods wherein the nucleic acid is introduced into the cells by naked DNA technology, adenovirus vector, adeno-associated virus vector, Epstein-Barr virus vector, Herpes virus vector, attenuated HIV vector, retroviral vectors, vaccinia virus vector, liposomes, antibody-coated liposomes, mechanical or electrical means. The above recited methods are merely served as examples for feasible means of introduction of the nucleic acid into cells. Other

099452-112404

- 8 -

methods known may be also be used in this invention.

5 In an embodiment of the above methods, the melanoma differentiation associated gene (*mda-7*) is linked to a regulatory element such that its expression is under the control of the regulatory element. In a still further embodiment, the regulatory element is inducible or constitutive. Inducible regulatory element like an
10 element such as promoter which can direct constitutive expression is also known in the art.

15 In a separate embodiment, the regulatory element is a tissue specific regulatory element. The expression of the *mda-7* gene will then be tissue-specific.

20 In another embodiment of the above-described methods, the cancer cell is characterized by the presence within the cancer cell of a defective tumor suppressor gene. The defective tumor suppressor gene includes, but is not limited to, a p53, a retinoblastoma (RB) or a p16^{ink4a} gene.

25 In an embodiment of the above-described methods, the cancer cell is characterized by the presence within the cancer cell of a dominant acting oncogene. Specifically, the dominant acting oncogene may be a Ha-ras, mutant p53 or human papilloma virus genes. The Ha-ras is a Harvey virus ras oncogene.

30 In an embodiment of the above methods, the nucleic acid comprises a vector. The vector includes, but is not limited to, an adenovirus vector, adeno-associated virus vector, Epstein-Barr virus vector, Herpes virus vector, attenuated HIV vector, retrovirus vector and vaccinia
35 virus vector. In a preferred embodiment, the adenovirus vector is a replication-defective adenovirus vector expressing *mda-7*, designated Ad.*mda-7* S. In another

05991452 "112101

embodiment, the adenovirus vector is a replication-competent adenovirus vector.

This invention also provides a method for reversing the cancerous phenotype of a cancer cell which comprises introducing the gene product of a melanoma differentiation associated gene (*mda-7*) into the cancerous cell so as to thereby reverse the cancerous phenotype of the cell.

10 This invention further provides a method for reversing the cancerous phenotype of a cancer cell in a subject which comprises introducing the gene product of a melanoma differentiation associated gene (*mda-7*) into the
15 subject's cancerous cell so as to thereby reverse the cancerous phenotype of the cell.

In an embodiment of the above-described methods, the cancer cell includes, but is not limited to, a breast, cervical, colon, prostate, nasopharyngeal, lung connective tissue or nervous system cell. The cancer cell further includes cells from glioblastoma multiforme, lymphomas and leukemia.

25 This invention also provides a pharmaceutical composition which comprises an amount of a nucleic acid comprising a melanoma differentiation associated gene (*mda-7*) effective to reverse the cancerous phenotype of a cancer cell and a pharmaceutically acceptable carrier.

30 As used herein, the term "pharmaceutically acceptable carrier" encompasses any of the standard pharmaceutical carriers. The pharmaceutical composition may be constituted into any form suitable for the mode of administration selected. Compositions suitable for oral
35 administration include solid forms, such as pills, capsules, granules, tablets, and powders, and liquid

forms, such as solutions, syrups, elixirs, and suspensions. Forms useful for parenteral administration include sterile solutions, emulsions, and suspensions.

15 This invention also provides a pharmaceutical composition comprising an amount of the gene product of a melanoma differentiation associated gene (*mda-7*) effective to reverse the cancerous phenotype of a cancer cell and a pharmaceutically acceptable carrier.

20 In an embodiment of the above-described methods, the cancer cell includes, but is not limited to, a breast, cervical, colon, prostate, nasopharyngeal, lung connective tissue and nervous system cells. The cancer

25 cell further includes cells from glioblastoma multiforme, lymphomas and leukemia.

This invention will be better understood from the Experimental Details which follow. However, one skilled in the art will readily appreciate that the specific methods and results discussed are merely illustrative of the invention as described more fully in the claims which follow thereafter.

- 11 -

Experimental Details

Cancer is a disease characterized by defects in growth control, and tumor cells often display abnormal patterns of cellular differentiation. The combination of recombinant human fibroblast interferon and the antileukemic agent mezerein corrects these abnormalities in cultured human melanoma cells resulting in irreversible growth arrest and terminal differentiation. Subtraction hybridization identifies a melanoma differentiation associated gene (*mda-7*) with elevated expression in growth arrested and terminally differentiated human melanoma cells. Colony formation decreases when *mda-7* is transfected into human tumor cells of diverse origin and with multiple genetic defects. In contrast, the effects of *mda-7* on growth and colony formation in transient transfection assays with normal cells, including human mammary epithelial, human skin fibroblast and rat embryo fibroblast, is quantitatively less than that found with cancer cells. Tumor cells expressing elevated *mda-7* display suppression in monolayer growth and anchorage independence. Infection with a recombinant type 5 adenovirus expressing antisense *mda-7* eliminates *mda-7* suppression of the *in vitro* growth and transformed phenotype. The ability of *mda-7* to suppress growth in cancer cells not expressing or containing defects in both the retinoblastoma (RB) and p53 genes indicates a lack of involvement of these critical tumor suppressor elements in mediating *mda-7*-induced growth inhibition. The lack of protein homology of *mda-7* with previously described growth suppressing genes and the differential effect of this gene on normal versus cancer cells suggests that *mda-7* may represent a new class of cancer growth suppressing genes with antitumor activity.

0591452.112101

- 12 -

MATERIALS AND METHODS

Cell Lines and Culture Conditions. Human carcinoma cell lines, including MCF-7 and T47D (breast), LS174T and SW480 (colorectal), HeLa (cervical), DU-145 (prostate), and HONE-1 (nasopharyngeal) (9,22-25), were grown in Dulbecco's modified Eagle's medium supplemented with 10% fetal bovine serum (DMEM-10) at 37°C in a 5% CO₂/95% air-humidified incubator. Additional human cell types including HBL-100 (normal mammary epithelial), H0-1 and C8161 (melanoma), GBM-18 and T98G (glioblastoma multiforme) and Saos-2 (human osteosarcoma) were maintained under similar conditions. Early passage normal human mammary epithelial cells (HMEC; passages 10-12) were obtained from Clonetics Corporation (San Diego, CA). HMEC cells were maintained in serum-free medium as described by Clonetics Corporation. CREF-Trans 6 (cloned Fischer rat embryo fibroblast) (9,26) and CREF Ha-ras (CREF cells transformed by the Ha-ras (T24) oncogene) (27) were cultured in DMEM-5. HeLa cl 1 is a hygromycin resistant (Hyg^R) Rous Sacroma virus RSV vector (pREP4) (Invitrogen) transformed HeLa clone. HeLa cl 2 is a Hyg^R *mda-7* expressing HeLa clone. HeLa cl 1 and HeLa cl 2 cells were constructed as described (12,21) and maintained in DMEM-10 containing 100 µg/ml of hygromycin. DU-145 cl 6 and DU-145 cl 7 cells contain a DEX-inducible *mda-7* gene (cloned in a pMAMneo vector) (Clontech) (21) and are maintained in DMEM-10 containing 200 µg/ml G418.

Subtraction Hybridization, Plasmids, Expression Vector Constructs, and Northern Hybridization. Identification and cloning of *mda-7* by subtraction hybridization was achieved as described (13). A full-length *mda-7* cDNA was isolated by screening a recombinant IFN-β plus MEZ-treated H0-1 cDNA library (13) and using the procedure of rapid amplification of cDNA ends as described (15). An *mda-7* cDNA fragment (nucleotide position 176-960) containing the open reading frame was

05991452-112101

- 13 -

amplified with PCR and cloned into pCRII™ (Invitrogen) by TA cloning. The orientation of the inserts in the vectors was determined by restriction mapping. The human cell expression constructs were made by cloning *Kpn I* - *Xho I* fragments from the PCR™ vectors into pREP4 vector (Invitrogen) downstream of a RSV promoter in a sense [*mda-7* (S)] or antisense [*mda-7* (AS)] orientation. Alternatively, the *mda-7* gene fragment was cloned into the pMAMneo (Clontech) vector in a sense and antisense orientation. RNA isolation and Northern blotting were performed as described (9,12,13,21).

Monolayer Growth, Anchorage-Independence and DNA-Transfection Assays. Monolayer and anchorage-independent growth assays were performed as previously described (8,12,26). To study the effect of *mda-7* on monolayer colony formation the vector [pREP4 (RSV)] containing no insert, *mda-7* (S) or *mda-7* (AS) expression constructs were transfected into the various cell types by the lipofectin method (GIBCO/BRL) and hygromycin resistant colony formation or cell growth in hygromycin was determined (12,21).

Construction of Antisense-*mda-7* Adenovirus Vector. The recombinant replication-defective Ad.*mda-7* (AS) was created in two steps. First, the coding sequence of the *mda-7* gene was cloned into a modified Ad expression vector pAd.CMV (28). This contains, in order, the first 355 bp from the left end of the Ad genome, the cytomegalovirus (CMV) immediate early promoter, DNA encoding splice donor and acceptor sites, cloning sites for the desired gene (in this case *mda-7*), DNA encoding a polyA signal sequence from the beta globin gene, and approximately 3 kbp of adenovirus sequence extending from within the E1B coding region. This arrangement allows high level expression of the cloned sequence by the CMV immediate early gene promoter, and appropriate RNA

0090452.12101

- 14 -

processing (28). The recombinant virus was created in vivo in 293 cells (29) by homologous recombination between *mda-7*-containing vector and plasmid JM17, which contains the whole of the Ad genome cloned into a modified version of pBR322 (30). JM17 gives rise to Ad genomes in vivo but they are too large to package. This constraint is relieved by recombination with the vector to create a packageable genome (30), containing the gene of choice. The recombinant virus is replication defective in human cells except 293 cells, which express adenovirus E1A and E1B. Following transfection of the two plasmids, infectious virus was recovered, the genomes were analyzed to confirm the recombinant structure, and then virus was plaque purified, all by standard procedures (31).

Peptide Antibody Production and Immunoprecipitation Analyses. Peptide antibodies were prepared against PSQENEMFSIRD as described (21). Logarithmically growing HeLa, HeLa cl 1 (Hyg^R pREP4 vector control HeLa clone), and HeLa cl 2 [pREP4-*mda-7* (S) transfected Hyg^R *mda-7* expressing HeLa clone] cells were either untreated or infected with 10 plaque forming units of control adenovirus (H5dl434) (32) or a recombinant adenovirus expressing *mda-7* (AS) [Ad.*mda-7* (AS)]. At various times after infection, cultures were starved of methionine for 1 hr at 37°C in methionine-free medium, cells were concentrated by pelleting and labeled for 4 hr at 37°C in 1 ml of the same medium with 100 µCi (1Ci=37GBq) of ³⁵S (NEN; Express ³⁵S). Immunoprecipitation analyses with 2 µg of MDA-7 peptide rabbit polyclonal antibody or actin monoclonal antibody (Oncogene Sciences) were performed as described (15,21).

EXPERIMENTAL RESULTS

Enhanced Growth Inhibitory Properties of *mda-7* in Human Cancer Cells and Ha-ras-Transformed Rat Embryo Fibroblast Cells. DNA transfection assays were performed to evaluate

- 15 -

the effect of elevated expression of *mda-7* on cell growth. When transfected into human cervical carcinoma (HeLa) cells, the *mda-7* (S) construct results in a 10- to 15-fold reduction in Hyg^R colonies in comparison with the pREP4 vector and *mda-7* (AS) construct transfected cultures (Fig. 1 and Table 1).

Table 1 Effect of *mda-7* on monolayer colony formation of human cancer, normal rat embryo fibroblast (CREF) and Ha-ras-transformed CREF cells.

Cell Type	RSV-Vector ^a	RSV- <i>mda-7</i> (S) ^b	RSV- <i>mda-7</i> (AS)
Human cancer cell lines ^c			
MCF-7 (Breast- Ca)	118 ± 24	42 ± 16 (3.5)	146 ± 20
T47D (Breast- Ca)	172 ± 9	44 ± 7 (4.2)	186 ± 28
HeLa (Cervix- Ca)	1571 ± 446	117±107 (15.2)	1771 ± 385
LS174T (Colorectal-Ca)	130 ± 14	30 ± 3 (5.4)	160 ± 15
HONE-1 (Nasopharyngeal-Ca)	219 ± 19	71±8 (3.5)	250 ± 19
DU-145 (Prostate-Ca)	174 ± 18	54 ± 8 (3.1)	166 ± 12
T98G (Glioblastoma)	99 ± 9	32 ± 4 (3.6)	115 ± 14
Saos-2 (Osteosarcoma)	126 ± 22	35 ± 6 (3.9)	138 ± 14
Rat embryo fibroblast			
CREF (normal rat embryo)	60 ± 10	35 ± 5 (1.7)	66 ± 7
CREF- <i>ras</i> (transformed)	147 ± 16	25 ± 4 (6.0)	151 ± 16

^a Logarithmically growing cells were seeded at 1 X 10⁶ per 100-mm plate and transfected with 10 µg of

000145-1201

- 16 -

- vector [pREP4 (RSV)] containing no insert, *mda-7* (S), or *mda-7* (AS). After 24 hr, cells were replated at approximately 2×10^5 cells per 100-mm plate in medium containing 100 μ g/ml of hygromycin. Medium was changed every 3 or 4 days and plates were fixed in formaldehyde and stained with Giemsa at day 14 or 21. Colonies containing 50 or more cells were enumerated. Values shown are the average Hyg^R colonies formed in four to five replicate plates \pm S.D.
- ^b Values in parentheses indicate fold-decrease in colony formation versus RSV-*mda-7* (AS) transfected cells.
- ^c MCF-7, T47D, HeLa, LS174T, DU-145 and HONE-1 are human carcinoma (Ca) cell lines isolated from the indicated anatomical site. T98G is a human glioblastoma multiforme cell line. CREF-ras is a Ha-ras (T24) oncogene transformed CREF clone.
- In addition to forming fewer colonies, *mda-7* (S) colonies are generally smaller in size than corresponding Hyg^R colonies resulting after transfection with the pREP4 vector or *mda-7* (AS) constructs (Fig. 1). When transfected into additional human cancer cell lines *mda-7* (S) constructs reduce Hyg^R colony formation by 3- to 10-fold (Table 1). These include human breast carcinoma (MCF-7 and T47D), colon carcinoma (LS174T and SW480), nasopharyngeal carcinoma (HONE-1), prostate carcinoma (DU-145), melanoma (H0-1 and C8161), glioblastoma multiforme (GBM-18 and T98G) and osteosarcoma (Saos-2). As observed with HeLa cells, the average sizes of Hyg^R colonies that form after transfection with *mda-7* (S) constructs are smaller than those formed following transfection with the empty pREP4 vector or *mda-7* (AS) constructs. These results demonstrate that *mda-7* is a potent growth suppressing gene when over-expressed in a wide spectrum of histologically distinct human cancers.

- 17 -

To determine if *mda-7* also inhibits the growth of normal cells and whether this effect is quantitatively similar to that observed with human cancer cells, transient DNA transfection assays were performed with passage 10 to 12 normal human mammary epithelial (HMEC) cells, the normal breast epithelial cell line HBL-100, normal human skin fibroblasts (passage 21) and a cloned normal rat embryo fibroblast cell line (CREF-Trans 6) (7,8). Since HMEC, HBL-100 and normal human skin fibroblasts do not form well-defined colonies at high frequencies, even when using a feeder-layer, the effect on total cell number after transfection with the different RSV constructs and growth for two and three weeks in hygromycin was determined. Using this approach, an approximate 1.1 to 1.6-fold decrease in HMEC, an approximate 1.1 to 1.2-fold decrease in HBL-100 and an approximate 1.3 to 2.1-fold decrease in normal human skin fibroblast cell number was observed (three independent experiments with each cell type) in *mda-7* (S) versus *mda-7* (AS) or pREP4 vector transfected normal cells, respectively. In contrast, using a similar experimental protocol with T47D human breast carcinoma cells, growth was inhibited following transfection with the *mda-7* (S) construct approximately 3.2 to 5.2-fold in comparison with vector- and antisense-transfected cells. In the case of CREF-Trans 6 cells, the difference in Hyg^R colony formation for six independent transfection assays between *mda-7* (S) versus *mda-7* (AS) and vector transfected cells ranged from 0.5 to 2.8-fold (Table 1). In contrast, transfection of *mda-7* (S) constructs into Ha-ras transformed CREF cells reduced colony formation by ~6 to 8-fold (Table 1). These results indicate that *mda-7* is quantitatively less effective in reducing growth and colony formation in normal human and normal rodent cells than in human cancer and Ha-ras-transformed rat embryo cells.

Effect of Stable and Inducible *mda-7* Expression and

009443E J2101

- 18 -

Antisense Inhibition of *mda-7* Expression on Cell Growth and the Transformed Phenotype. To determine the reason for low frequency HeLa cell survival after transfection with the *mda-7* (S) gene, ten independent Hyg^R colonies were isolated following transfection with the *mda-7* (S) construct. Of the 10 clones analyzed by Northern blotting for *mda-7* expression, 7 clones did not express detectable *mda-7* mRNA, 2 clones expressed low levels of *mda-7* mRNA and 1 clone (designated HeLa cl 2) displayed high levels of *mda-7* mRNA. In contrast, all of the clones displayed comparable levels of Hyg^R and glyceraldehyde 3-phosphate dehydrogenase (GAPDH) gene expression. When compared with parental HeLa cells or an pREP4 vector HeLa clone (designated HeLa cl 1), HeLa cl 2 (*mda-7* expressing) cells grew at a reduced rate (Fig. 2). When grown in agar, uncloned HeLa and HeLa cl 1 cells grew with approximately 42% efficiency, whereas HeLa cl 2 (*mda-7* expressing) cells grew with approximately 25% efficiency and the average sizes of colonies were smaller than observed with parental HeLa and pREP4 vector HeLa cl 1 cells. These results indicate that HeLa survival after transfection with *mda-7* results primarily from the lack of or low levels of *mda-7* expression. However, in HeLa cells that stably express elevated *mda-7*, growth in monolayer culture and anchorage-independence are reduced.

To determine if the reduction in *in vitro* growth and transformation suppression found in HeLa cl 2 (*mda-7* expressing) are a direct consequence of *mda-7* expression, an antisense strategy was used to directly inhibit *mda-7* expression. A recombinant Ad5 vector containing the *mda-7* gene cloned in an antisense orientation [Ad.*mda-7* (AS)] was constructed. Infection of HeLa cl 2 (*mda-7* expressing), but not HeLa cl 1 (pREP4 vector, non-*mda-7* expressing) or parental HeLa, with Ad.*mda-7* (AS) increases growth rate and agar cloning efficiency (from approximately 25 to approximately 44%) (Fig. 2). In

- 19 -

contrast, the control mutant Ad5 vector (H5dl434), not containing the *mda-7* gene, does not affect monolayer or agar growth of parental HeLa, HeLa cl 1 or HeLa cl 2 cells (data not shown).

5

Using *mda-7*-specific peptide antibodies produced in rabbits and immunoprecipitation analyses, the HeLa cl 2 (*mda-7* expressing) cells contain elevated levels of the MDA-7 approximately 24 kDa protein and a high molecular weight complexing (HMC) protein of approximately 90 to 110 kDa (Fig. 3). Infection with Ad.*mda-7* (AS), but not the H5dl434 control non-*mda-7* expressing virus, results in a temporal decrease in both the ~24 kDa MDA-7 protein and the HMC protein (21) (Fig. 3). Reduced levels of both proteins are seen by 48 hr and remain suppressed over a 96 hr period after infection with Ad.*mda-7* (AS). In contrast, actin levels remain unaltered following viral infection. These findings indicate that antisense inhibition of MDA-7 protein expression in HeLa cl 2 (*mda-7* expressing) can directly extinguish *mda-7* induced growth suppression and inhibition in anchorage-independent growth.

To confirm the suppressive effect of *mda-7* on cell growth, DU-145 human prostate cancer cells were engineered to express a DEX-inducible *mda-7* gene. When DU-145 cl 6 or cl 7 cells [containing a DEX-inducible *mda-7* (S) gene], but not parental DU-145 cells, are grown for 24 to 96 hr in the presence of 10^{-6} M DEX, *mda-7* mRNA and protein (including the HMC protein) are induced (Fig. 4). In contrast, DEX does not alter neomycin resistance (Neo^R) gene expression in DU-145 cl 6 and cl 7 cells or GAPDH expression in any of the cells tested (Fig. 4). Induction of *mda-7* expression in DU-145 cl 6 and cl 7 cells by growth in 10^{-6} M DEX results in approximately 50% reduction in cell number after 96 hr versus growth in the absence of DEX. In contrast, no significant growth

0991450.1866

- 20 -

inhibition occurs when parental DU-145 or pMAMneo vector transformed DU-145 cells are grown for 96 hr in medium containing 10^{-6} M DEX (data not shown). These data indicate that ectopic expression of *mda-7* can directly alter cell growth in prostate cancer cells.

EXPERIMENTAL DISCUSSION

Subtraction hybridization identified *mda* genes with elevated expression in growth arrested and terminally differentiated human melanoma cells (13,14,21). Determining the function of these *mda* genes will be paramount in defining the molecular basis of growth control and terminal differentiation in human melanoma and other cell types. The *mda-7* gene (14,21) is now shown to be a ubiquitous growth suppressing gene when transiently or stably expressed in a wide array of human cancer cell lines. This finding extends previous observations indicating growth inhibitory properties of the MDA-7 protein in human melanoma cells (21). In contrast to its effects on cancer cells, transfection of *mda-7* into normal human mammary epithelial, normal human skin fibroblast and normal rat embryo fibroblast cells results in quantitatively less growth suppression. Like another *mda* gene, *mda-6* (p21), *mda-7* expression is also inversely correlated with melanoma progression, with elevated levels of both *mda-6* (p21) and *mda-7* present in normal human melanocytes relative to metastatic human melanoma cells (14-16,21). Since normal melanocytes still retain proliferative capacity, although at a reduced rate relative to melanoma cells, it is possible that both *mda-6* (p21) and *mda-7* function as negative regulators of the progression phenotype in melanocyte/melanoma lineage cells (14-16,21). Moreover, the elevated expression of both *mda-6* (p21) and *mda-7* in terminally differentiated and irreversibly growth arrested human melanoma cells, suggests that these genes may also be important regulators of the terminal differentiation phenotype

- 21 -

(13-16,21).

5 The mechanism by which *mda-7* elicits its growth suppressive effects on human cancer cells is not presently known. The structure of *mda-7* does not provide insight into potential function, since no sequence motifs are present that would suggest a potential mode of action. The effect of *mda-7* on cell growth can be distinguished from the extensively studied tumor suppressor gene p53 (33,34). Transient expression of p53 in the mutant p53 containing T47D human breast carcinoma cell line results in growth suppression, whereas transfection of a wild-type p53 gene into the wild-type p53 containing MCF-7 human breast carcinoma cell line does not induce growth inhibition (34). In contrast, *mda-7* induces similar growth suppression in both T47D and MCF-7 cells (Table 1). Growth inhibition by *mda-7* can also be disassociated from that observed with the retinoblastoma gene (pRb), the pRb-associated p107 gene and the putative tumor suppressor gene p16^{ink4} (25,35). Overexpression of pRb and p107 inhibit cellular proliferation in specific cell types and in a cell cycle-dependent manner (35-37). Transfection of pRb or p107 into the human glioblastoma cell line T98G that contains an apparently normal RB gene (25) does not induce growth suppression (35,37), whereas transient *mda-7* (S) expression reduces T98G colony formation (Table 1). At the present time, the growth inhibitory effect of *mda-7* cannot be distinguished from growth suppression induced by the RB family member p130/pRb2, which also inhibits proliferation in T98G cells (25). The p16^{ink4} gene induces growth arrest in cells containing a functional RB gene (35,37), whereas *mda-7* growth suppression occurs in cells containing normal, abnormal or non-functional RB genes. Transfection of *mda-7* into the DU-145 human prostate carcinoma cell line that contains a mutated RB gene (38) and Saos-2 human osteosarcoma cells that do not

009445E-412104

- 22 -

express RB (or wild-type p53) results in an inhibition in colony formation (Table 1). Similarly, induction of *mda-7* expression in stable DEX-inducible *mda-7* transformed DU-145 clones results in growth suppression. These findings indicate a lack of dependence on a functional RB gene for growth inhibition by *mda-7*. Taken together these studies demonstrate that the inhibitory effect of *mda-7* occurs by a mechanism that is distinct from the mode of action of the two most extensively studied tumor suppressor genes, p53 and pRb, and the putative tumor suppressor gene p16^{ink4}.

Several genes have been identified that display elevated expression as a function of growth arrest or DNA damage in mammalian cells (39,40). Three growth arrest and DNA damage inducible (*gadd*) genes, *gadd45*, *gadd153* and *gadd34*, the closely related myeloid differentiation primary response (*MyD118*) gene (41) and the wild-type p53 inhibiting gene *mdm-2* (42) are upregulated in cells by treatment with the DNA damaging agent methyl methanesulfonate (MMS) (40). The *gadd45* and growth arrest-specific gene (*gas1*) (43,44) are induced by maintaining cells at confluence, serum-starving cells or growing cells in low serum (40,43,44). In contrast, *mda-7* mRNA expression is not induced in human melanoma cells following treatment with methyl methane sulfonate (MMS) or after maintaining cells at confluence (21). Moreover, only a small increase in *mda-7* mRNA expression occurs in H0-1 human melanoma cells following growth in serum-free medium for 96 hr (21). The difference in regulation of *mda-7* versus the *gadd*, *MyD118* and *gas-1* genes indicates that *mda-7* may represent a new class of growth arresting genes.

In summary, a negative growth regulator, *mda-7*, is described that induces growth suppression in human cancer cells containing both normal and mutated p53 and RB

09945-120

- 23 -

genes. Characterization of the genomic structure of *mda-7* will be important in determining if this gene normally functions as a tumor suppressor gene and whether alterations are present in this gene in tumor versus normal cells. Identification of the promoter region of *mda-7* will also permit an analysis of the mechanism by which this gene is differentially expressed and inducible by IFN- β plus MEZ in specific cell types. Of potential importance and warranting expanded studies, is the finding that *mda-7* is more growth inhibitory toward cancer and transformed cells than normal cells. In this context, *mda-7* could prove useful as part of a gene-based interventional strategy for cancer therapy, in an analogous manner as the wild-type p53 gene is currently being tested for efficacy in the therapy of specific human malignancies.

00001452 112101

References

1. Fisher, P.B. (1984) in *Tumor Promotion and Cocarcinogenesis in Vitro: Mechanisms of Tumor Promotion*, ed. Slaga, T.J., (CRC, Boca Raton, FL),
5 vol. 3, pp. 57-123.
2. Bishop, J.M. (1991). *Cell* 64, 235-248.
3. Knudson, A.G. (1991). *Proc. Natl. Acad. Sci. USA* 90,
10914-10921.
4. MacLachlan, T.K., Sang, N. & Giordano, A. (1995).
10 *Crit. Rev. Eukaryotic Gene Express.* 5, 127-156.
5. Sang, N., Baldi, A. & Giordano, A. (1995). *Mol. Cell. Differ.* 3, 1-29.
6. Barbacid, M. (1987). *Annu. Rev. Biochem.* 56,
779-827.
- 15 7. Bos, J. (1989). *Cancer Res.* 49, 4682-4689.
8. Su, Z.-z., Olsson, C.A., Zimmer, S.G. & Fisher, P.B. (1992). *Anticancer Res.* 12, 297-304.
9. Shen, R., Su, Z.-z., Olsson, C. A. & Fisher, P. B. (1995). *Proc. Natl. Acad. Sci. USA* 92, 6778-6782.
- 20 10. Noda, M., Kitayama, H., Matsuzaki, T., Sugimoto, Y., Okayama, H., Bassin, R. H. & Ikawa, Y. (1988) *Proc. Natl. Acad. Sci. USA* 86, 162-166.
11. Kitayama, H., Sugimoto, Y., Masuzaki, T., Ikawa, Y. & Noda, M. (1989). *Cell* 56, 77-84.
- 25 12. Lin, J., Su, Z.-z., Grunberger, D., Zimmer, S. G. & Fisher, P. B. (1994). *Int. J. Oncol.* 5, 5-15.
13. Jiang, H. & Fisher, P. B. (1993). *Mol. Cell. Differ.* 1, 285-299.
14. Jiang, H., Lin, J. & Fisher, P. B. (1994). *Mol. Cell. Differ.* 2, 221-239.
- 30 15. Jiang, H., Lin, J., Su, Z.-z., Herlyn, M., Kerbel, R. S., Weissman, B. E., Welch, D. R., & Fisher, P. B. (1995) *Oncogene* 10, 1855-1864.
16. Jiang, H., Lin, J., Su, Z.-z. & Fisher, P. B. (1996). *Mol. Cell. Differ.* 4, 67-89.
- 35 17. Xiong, Y., Hannon, G. J., Zhang, H., Casso, D., Kobayashi, R. & Beach, D. (1993). *Nature (London)*

090116 110 120 130 140 150 160 170 180 190 200 210 220 230 240 250 260 270 280 290 300 310 320 330 340 350 360 370 380 390 400 410 420 430 440 450 460 470 480 490 500 510 520 530 540 550 560 570 580 590 600 610 620 630 640 650 660 670 680 690 700 710 720 730 740 750 760 770 780 790 800 810 820 830 840 850 860 870 880 890 900 910 920 930 940 950 960 970 980 990

- 25 -

366, 701-704.

18. El-Deiry, W. S., Tokino, T., Velculescu, V. E.,
Levy, D. B., Parsons, R., Trent, J. M., Lin, D.,
Mercer, W. E., Kinzler, K. W. & Vogelstein, B.
5 (1993). *Cell* 75, 817-825.
19. Harper, J. W., Adami, G. R., Wei, N., Keyomarsi, K. &
Elledge, S. J. (1993). *Cell* 75, 805-816.
20. Noda, A., Ning, Y., Venable, S. F., Pereira-Smith,
O. M. & Smith, J. R. (1994). *Exp. Cell Res.* 211,
10 90-98.
21. Jiang, H., Lin, J. J., Su, Z.-z., Goldstein, N. I.
& Fisher, P. B. (1995). *Oncogene* 11, 2477-2486.
22. Leon, J. A., Mesa-Tejada, R., Gutierrez, M. C.,
Estabrook, A., Greiner, J. W., Schlom, J. & Fisher,
15 P. B. (1989). *Anticancer Res.* 9, 1639-1648.
23. Kantor, J., Tran, R., Greiner, G., Pestka, S.,
Fisher, P. B., Shively, J. E. & Schlom, J. (1989).
Cancer Res. 49, 2651-2655.
24. Leon, J. A., Gutierrez, M. C., Jiang, H., Estabrook,
20 A., Waxman, S. & Fisher, P. B. (1992). *Cancer
Immunol. Immunother.* 35, 315-324.
25. Claudio, P. P., Howard, C. M., Baldi, A., De Luca,
A., Fu, Y., Condorelli, G., Sun, Y., Colburn, N.,
Calabretta, B. & Giordano, A. (1994). *Cancer Res.*
25 54, 5556-5560.
26. Fisher, P.B., Babiss, L. E., Weinstein, I. B. &
Ginsberg, H.S. (1982). *Proc. Natl. Acad. Sci. U.S.A.*
79, 3527-3531.
27. Boylon, J., Shih, T., Fisher, P. B. & Zimmer, S. G.
30 (1992). *Mol. Carcinog.* 5, 118-128.
28. Falck-Pedersen, E., Heinflink, M., Alvira, M.,
Nussenzweig, D.R. & Gershengorn, M.C. (1994). *Mol.
Pharmacol.* 45, 684-689.
29. Graham, F.L., Smiley, J., Russell, W.C. & Nairn, R.
35 (1977). *J. Gen. Virol.* 36, 59-72.
30. McGrory, W.J., Bautista, D.S. & Graham, F.L. (1988).
Virology 163, 614-617.

096452 12104

- 26 -

31. Volkert, F.C. & Young, C.S.H. (1983). *Virology* **125**, 175-193.
32. Grodzicker, T. & Klessig, D. (1980). *Cell* **21**, 453-463.
- 5 33. Baker, S.J., Markowitz, S., Fearon, E.R., Wilson, J.K.V. & Vogelstein, B. (1990). *Science* **249**, 912-915.
34. Mercer, W.E. (1992). *Crit. Rev. Eukaryotic Gene Expression* **2**, 251-263.
- 10 35. Medema R. H., Herrera, R. E., Lam F. & Weinberg, R. A. (1995). *Proc. Natl. Acad. Sci. USA* **92**, 6289-6293.
36. Grana, X. & Reddy, E. P. (1995). *Oncogene* **11**, 211-220.
37. Zhu, L., van den Heuvel, S., Helin, K., Fattaey, A., Ewen, M., Livingston, D., Dyson, N. & Harlow, E. (1993). *Genes & Dev.* **7**, 1111-1125.
- 15 38. Bookstein, R., Shew, J. Y., Chen, P. L., Scully, P. & Lee, W. H. (1990). *Science* **247**, 712-715.
39. Fornace, A. J., Jr., Alamo, I. J., Jr. & Hollander, M. C. (1988). *Proc. Natl. Acad. Sci. USA* **85**, 8800-8804.
- 20 40. Zhan, Q., Lord, K.A., Alamo, I., Jr., Hollander, M.C., Carrier, F., Ron, D., Kohn, K. W., Hoffman, B., Liebermann, D.A. & Fornace, A. J., Jr. (1994). *Mol. Cell. Biol.* **14**, 2361-2371.
- 25 41. Abdollahi, A., Lord, A., Hoffman-Liebermann, B. & Liebermann, D. (1991). *Oncogene* **6**, 165-167.
42. Momand, J., Zambetti, G. P., Olson, D. C., George, D. & Levine, A. J. (1992). *Cell* **69**, 1237-1245.
- 30 43. Schneider, C., King, R.M. & Philipson, L. (1988). *Cell* **54**, 787-793.
44. Del Sal, G., Ruaro, M.E., Philipson, L. & Schneider, C. (1992). *Cell* **70**, 595-607.
45. Sambrook, J., et al. (1989) Molecular Cloning: a laboratory manual. Second Edition. Cold Spring Harbor Laboratory Press (1989).
- 35

 09945-120
 T024T660

- 27 -

Second Series of Experiments

Melanoma Differentiation Associated Gene-7 (*mda-7*) in a Recombinant Adenovirus Inhibits the Growth of Established Human Tumors in Nude Mice

5

Previous studies document that ectopic expression of *mda-7* in human tumor cells of diverse origins inhibits growth, as documented by a decrease in colony formation in monolayer culture (Jiang et al., PNAS, 93: 9160-9165, 1996). In contrast, *mda-7* does not significantly alter the growth of normal human epithelial or fibroblast cells. These observations support the hypothesis that *mda-7* is a ubiquitous cancer growth suppressor gene.

10

15 The ability of *mda-7* to selectively inhibit cancer cell growth suggests that this gene might provide therapeutic benefits in the treatment of human cancers. To explore this possibility a replication-defective adenovirus expressing *mda-7* has been generated. The protocols were similar to those used to construct an adenovirus expressing antisense *mda-7*, Ad.*mda-7* AS (Jiang et al., PNAS, 93: 9160-9165, 1996). The recombinant replication-defective Ad.*mda-7* S was produced in two steps. First, the *mda-7* gene was cloned in a sense orientation into a modified Ad expression vector pAd.CMV. This virus contains, in order, the first 355 bp from the left end of the Ad genome, the cytomegalovirus (CMV) immediate early promoter, DNA encoding a poly A signal sequence from the beta globin gene, and approximately 3 kbp of adenovirus sequence extending from within the E1B coding region. This arrangement allows high level expression of the cloned sequence by the CMV immediate early gene promoter, and appropriate RNA processing. The recombinant virus was created *in vivo* in 293 cells by homologous recombination between *mda-7* containing vector and JM17, which contains the whole of the Ad genome cloned into a modified version of pBR322. JM17 gives rise

20

25

30

35

009415E.12.10.1

- 28 -

to Ad genomes *in vivo* but they are too large to package. This constraint is relieved by recombination with the vector to create a packageable genome, containing the gene of choice. The recombinant virus is replication
5 defective in human cells except 293, which express adenovirus E1A and E1B. Following transfection of the two plasmids, infectious virus was recovered, the genomes were analyzed to confirm the recombinant structure, and then virus was plaque purified, all by standard
10 procedures.

As observed with transfection with *mda-7*, infection of diverse human cancer cell lines, but not normal cell lines, with Ad.*mda-7* S inhibited growth. These results
15 demonstrate that this virus retains properties observed with the *mda-7* plasmid construct. In many cancer cells, including breast carcinoma (MCF-7 and T47D), glioblastoma (GBM-18 and T98G) and melanoma (H0-1 and C8161), infection with Ad.*mda-7* S resulted in the induction of
20 programmed cell death (apoptosis). This effect was not elicited in normal cells even after infection with high multiplicities of infection (100 pfu/cell) with Ad.*mda-7* S. In other cancer cell types, growth suppression (as indicated by a suppression in colony formation in
25 monolayer culture) was apparent without signs of apoptosis, as indicated by nuclear morphology changes, formation of nucleosomal ladders or a positive TUNEL reaction. These results indicate that the Ad.*mda-7* S virus can selectively inhibit the growth of human cancer
30 cells *in vitro*. Moreover, in specific cancer cell types growth suppression correlates with induction of apoptosis. These observations suggest that inhibition in cancer growth induced by *mda-7* can occur by multiple pathways.

35

Nude mouse human tumor xenograft models were used to determine if Ad.*mda-7* S can inhibit the growth of human

099445-1-101

DRAFT

DRAFT

DRAFT